

## ***Interactive comment on* “Storm Tide Amplification and Habitat Changes due to Urbanization of a Lagoonal Estuary” by Philip M. Orton et al.**

**Carmine Donatelli (Referee)**

c.donatelli@liverpool.ac.uk

Received and published: 20 April 2020

I have read with great interest the paper entitled: ‘Storm Tide Amplification and Habitat Changes due to Urbanization of a Lagoonal Estuary’, by Orton et al.

The manuscript is very well written and organized, the used methodology is robust and supported by the clearness of the presented data. Previous works are referenced appropriately.

I think this study represents an important contribution to our understanding of the mechanisms by which geomorphic changes alter the physical response of back-barrier estuaries to tides and low frequency actions. I recommend publication with minor revisions.

My minor comments/suggestions are listed below.

C1

[Printer-friendly version](#)

[Discussion paper](#)



Pages 10-11: 'The most dramatic land cover change is from large areas of fringing wetlands (light blue) to urbanized areas (red), but also the center of the bay has shifted from marshes to open waters (1 dark blue)'.

The influence of marsh loss on water levels is strongly related to the setting: marsh loss associated with lateral erosion vs. marsh loss associated with reclamation projects. Reductions in the basin planform area (wetland reclamation) increase water levels, while marsh retreat due to lateral erosion may have the opposite effect (in agreement with the analytical model of Keulegan, 1967 and 3D numerical modelling investigations, Picado et al., 2010; Donatelli et al., 2018). I would highlight/expand this point in the text.

Page 15: 'It was previously established that the bay's tide ranges have grown substantially (Swanson and Wilson, 2008), and we find similar results. Averaging high and low waters for daytime minima and maxima in 1878 over 37 days gives an observed tide range of 1.35 m, while observations for the entire year 2015 show a tide range of 1.73 m. This increase of 28% is smaller than the prior estimate of the tide range change from 1899 to 2000 from Swanson and Wilson (2007), which was 1.16 m to 1.64 m or 41%. However, the 1878 measurements are for a location at mid-bay (Holland House), whereas the 1899 measurements are for the easternmost end of the bay (Inwood or Norton Point), where tide attenuation (e.g. due to narrow, shallow channels and wetlands) was likely more pronounced.'

I would move these lines into the Discussion section.

Page 16 (lines 11-14): I would expand these lines. Are the results for Jamaica Bay translatable to other systems? What systems?

Page 18 (lines 1-19): I would move the description of the idealized numerical experiments into the Methods section.

Page 21: I would mention Englebright [1975] and Harting et al., [2002], who show how

[Printer-friendly version](#)[Discussion paper](#)

inlet modifications have reduced the movement of offshore sediments into the back-barrier basin.

---

Interactive comment on Nat. Hazards Earth Syst. Sci. Discuss., <https://doi.org/10.5194/nhess-2019-343>, 2020.

[Printer-friendly version](#)

[Discussion paper](#)

